

---

## The Possible Benefits of NanoTek Chitosan, a Dietary Supplement

- May support enhanced detoxification of Herxheimer's reactions\*
- Promotes the growth of most species of friendly bifidobacteria and lactobacilli
- Helps prevent lipid peroxidation, and promotes the activity of antioxidant enzymes\*
- Potentially protects against toxins, including mercury, carbon tetrachloride, and dioxins\*
- Potentially provides support for normal blood sugar regulation\*
- May have immunostimulative effects\*

---

## Description

NanoTek Chitosan contains chitosan oligosaccharides as well as a small amount of chitosan. Chitosan oligosaccharide has a much smaller molecular size than chitosan, and can absorb into the bloodstream. Both are derived from chitin, which is found in the exoskeletons of crustaceans such as shrimp, lobster or crab.

Chitosan is an aminopolysaccharide that is chemically similar to cellulose. Chitosan oligosaccharides (CO) are manufactured from chitosan by an enzymatic process, resulting in a much smaller molecular size that is more easily absorbed by the body. Chitosan itself is more of a fiber and contributes to detoxification in the GI tract, whereas CO supports detoxification in blood and other tissues.\*

In a small clinical trial of people being treated for Borrelia and experiencing Herxheimer's reactions, NanoTek Chitosan improved Herxheimer's symptoms by the end of one week.\* In subjects with pain, 100% reported significant improvement (in half of these, the pain completely resolved), and of those with sleep

disturbances, 75% reported sleeping more deeply and waking up more well rested.\*

The research on chitosan oligosaccharides is extensive, and shows it has great potential to benefit health. Chitosan oligosaccharides can promote the growth of friendly bifidobacteria and lactobacillus. Unlike fructooligosaccharides (FOS), which promote the growth of only three probiotic strains, chitosan oligosaccharide supports almost all bifido- and lactobacillus species.

Chitosan oligosaccharide has been shown to protect the liver from damage by carbon tetrachloride in mice, and to protect against mercury toxicity.\* It can promote the activity of antioxidant enzymes and help prevent lipid peroxidation, helping to protect mice poisoned with dioxin. It has also shown potential to support microbial balancing and other immune functions, and may help support normal blood sugar levels.\*

Investigated and utilized for many years by Akira Matsunaga, M.D., Ph.D., chitosan

itself is considered a superior health substance in Japan, because it has a broad effect on all of the body's systems. Dr. Matsunaga found that it strengthened those who were weak, made healthy patients healthier, and improved common daily complaints and the quality of life.\* He found that chitosan did not target only one organ, but effectively supported the functioning of all of the body's systems and organs.\*

contains more in-depth discussions of both chitosan and chitosan oligosaccharide, at <http://www.allergyresearchgroup.com/July-2007-Focus-Newsletter-sp-41.html>.

Both chitosan and chitosan oligosaccharide are extremely safe. Even at a chitosan oligosaccharide intake of 2000 mg/kg in rats, which extrapolates to more than 135,000 mg per day for an average-weight adult human, researchers could not find a potential adverse effect or toxicity.

Our July 2007 ARG Focus Newsletter

---

Serving Size: 2 Vegetarian Capsules

Servings Per Container: 30

**Amount Per Serving:**

Proprietary Blend

1200 mg

Chitosan Oligosaccharide and Chitosan (shrimp, crab, lobster)

Other ingredients: Hydroxypropyl methylcellulose, cellulose, L-leucine.

**Suggested Use:** As a dietary supplement, 1 capsule three times daily or 2 capsules two times daily on an empty stomach, or as directed by a healthcare practitioner.

**WARNING:** Avoid this product if you are allergic to shellfish, pregnant or lactating. If used long-term, additional supplementation with fat soluble vitamins and essential fatty acids is recommended.

---

**References**

Lee HW, Park YS, Jung JS, Shin WS. Chitosan oligosaccharides, dp 2-8, have prebiotic effect on the Bifidobacterium bifidum and Lactobacillus sp. Anaerobe. 2002 Dec;8(6):319-24.

Yan Y, Wanshun L, Baoqin H, Bing L, Chenwei F. Protective effects of chitosan oligosaccharide and its derivatives against carbon tetrachloride-induced liver damage in mice. Hepatol Res. 2006 Jul;35(3):178-84. Epub 2006 May 26.

Yoon HJ, Park HS, Bom HS, Roh YB, Kim JS, Kim YH. Chitosan oligosaccharide inhibits 203HgCl<sub>2</sub>-induced genotoxicity in mice: micronuclei occurrence and chromosomal aberration. Arch Pharm Res. 2005 Sep;28(9):1079-85.

Lee HW, Park YS, Choi JW, Yi SY, Shin WS. Antidiabetic effects of chitosan oligosaccharides in neonatal streptozotocin-induced noninsulin-dependent diabetes mellitus in rats. Biol Pharm Bull. 2003 Aug;26(8):1100-3.

Shon YH, Park IK, Moon IS, Chang HW, Park IK, Nam KS.

Effect of chitosan oligosaccharide on 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced oxidative stress in mice. Biol Pharm Bull. 2002 Sep;25(9):1161-4.

Moon JS, Kim HK, Koo HC, Joo YS, Nam HM, Park YH, Kang MI. The antibacterial and immunostimulative effect of chitosan-oligosaccharides against infection by Staphylococcus aureus isolated from bovine mastitis. Appl Microbiol Biotechnol. 2007 Mar 15; [Epub ahead of print]

Kim SK, Park PJ, Yang HP, Han SS. Subacute toxicity of chitosan oligosaccharide in Sprague-Dawley rats. Arzneimittelforschung. 2001 Sep;51(9):769-74.

Unpublished study, Steven W. Hines N.D., N.E., Hope Clinics International.

Matsunaga A. Chitosan, The Ultimate Health Builder. Vintage Press, NY, NY, 1998.



Allergy Research Group®  
2300 North Loop Road, Alameda, CA 94502  
Phone: 800-545-9960 or 510-263-2000  
Fax: 800-688-7426 or 510-263-2100  
[www.AllergyResearchGroup.com](http://www.AllergyResearchGroup.com)

\*THESE STATEMENTS HAVE NOT BEEN EVALUATED BY THE FOOD AND DRUG ADMINISTRATION. THIS PRODUCT IS NOT INTENDED TO DIAGNOSE, TREAT, CURE, OR PREVENT ANY DISEASE.